arXiv:1303.7370v1 [math.CA] 29 Mar 2013

HERMITE-HADAMARD TYPE INEQUALITIES FOR MAPPINGS WHOSE DERIVATIVES ARE $s-$CONVEX IN THE SECOND SENSE VIA FRACTIONAL INTEGRALS

ERHAN SET$^{\blacktriangle}$, M. EMIN OZDEMIR$^{\bigstar}$, M. ZEKI SARIKAYA$^{\blacksquare}$, AND FILIZ KARAKOÇ$^{\clubsuit}$

Abstract. In this paper we establish Hermite-Hadamard type inequalities for mappings whose derivatives are $s-$convex in the second sense and concave.

1. **Introduction**

Let f:$I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ be a convex function defined on the interval I of real numbers and $a,b\in I$ with $a<b$. Then

$$f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int\limits_{a}^{b}f(x)dx\leq\frac{f(a)+f(b)}{2} \tag{1.1}$$

is known that the Hermite-Hadamard inequality for convex function. Both inequalities hold in the reserved direction if f is concave. We note that Hadamard's inequality may be regarded as a refinement of the concept of convexity and it follows easily from Jensen's inequality. Hadamard's inequality for convex functions has received renewed attention in recent years and a remarkable variety of refinements and generalizations have been found; see, for example see ([5]-[17]).

Definition 1. *([4]) A function $f:[0,\infty)\rightarrow\mathbb{R}$ is said to be $s-$convex in the second sense if*

$$f(\lambda x+(1-\lambda)y)\leq\lambda^{s}f(x)+(1-\lambda)^{s}f(y)$$

for all $x,y\in[0,\infty),\lambda\in[0,1]$ and for some fixed $s\in(0,1]$. This class of $s-$convex functions is usually denoted by K_{s}^{2}.

In ([3]) Dragomir and Fitzpatrick proved a variant of Hadamard's ineqality which holds for $s-$convex functions in the second sense:

Theorem 1. *Suppose that $f:[0,\infty)\rightarrow[0,\infty)$ is an $s-$convex function in the second sense ,where $s\in(0,1)$ and let $a,b\in[0,\infty),a<b$. If $f'\in L^{1}([a,b])$, then the following inequalities hold:*

$$2^{s-1}f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int_{a}^{b}f(x)dx\leq\frac{f(a)+f(b)}{2} \tag{1.2}$$

The constant $k=\frac{1}{s+1}$ is the best possible in the second ineqality in (1.2)

2000 *Mathematics Subject Classification.* 26A33, 26A51, 26D07, 26D10, 26D15.

Key words and phrases. Hermite-Hadamard type inequality, $s-$convex function, Riemann-Liouville fractional integral.

The following results are proved by M.I.Bhatti et al. (see [2]).

Theorem 2. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a twice differentiable function on I° such that $|f''|$ is convex function on I. Suppose that $a, b \in I^{\circ}$ with $a < b$ and $f'' \in L[a,b]$, then the following ineqality for fractional integrals with $\alpha > 0$ holds:*

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha} f(b) + J_{b^-}^{\alpha} f(a)\right]\right| \\ &\leq \frac{(b-a)^2}{\alpha+1}\beta(2,\alpha+1)\left[\frac{|f''(a)|+|f''(b)|}{2}\right] \end{aligned} \tag{1.3}$$

where β is Euler Beta function.

Theorem 3. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a twice differentiable function on I°. Assume that $p \in \mathbb{R}, p > 1$ such that $|f''|^{\frac{p}{p-1}}$ is convex function on I. Suppose that $a, b \in I^{\circ}$ with $a < b$ and $f'' \in L[a,b]$, then the following ineqality for fractional integrals holds:*

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha} f(b) + J_{b^-}^{\alpha} f(a)\right]\right| \\ &\leq \frac{(b-a)^2}{\alpha+1}\beta^{\frac{1}{p}}(p+1,\alpha p+1)\left(\frac{|f''(a)|^q+|f''(b)|^q}{2}\right)^{\frac{1}{q}} \end{aligned} \tag{1.4}$$

where β is Euler Beta function.

Theorem 4. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a twice differentiable function on I°. Assume that $q \geq 1$ such that $|f''|^q$ is convex function on I.Suppose that $a, b \in I^{\circ}$ with $a < b$ and $f'' \in L[a,b]$,then the following ineqality for fractional integrals holds:*

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha} f(b) + J_{b^-}^{\alpha} f(a)\right]\right| \\ &\leq \frac{\alpha(b-a)^2}{4(\alpha+1)(\alpha+2)}\left[\begin{array}{c} \left(\frac{2\alpha+4}{3\alpha+9}|f''(a)|^q + \frac{\alpha+5}{3\alpha+9}|f''(b)|^q\right)^{\frac{1}{q}} \\ +\left(\frac{\alpha+5}{3\alpha+9}|f''(a)|^q + \frac{2\alpha+4}{3\alpha+9}|f''(b)|^q\right)^{\frac{1}{q}} \end{array}\right] \end{aligned} \tag{1.5}$$

Theorem 5. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a twice differentiable function on I°. Assume that $p \in \mathbb{R}, p > 1$ with $q = \frac{p}{p-1}$ such that $|f''|^q$ is concave function on I.Suppose that $a, b \in I^{\circ}$ with $a < b$ and $f'' \in L[a,b]$,then the following ineqality for fractional integrals holds:*

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha} f(b) + J_{b^-}^{\alpha} f(a)\right]\right| \\ &\leq \frac{(b-a)^2}{\alpha+1}\beta^{\frac{1}{p}}(p+1,\alpha p+1)\left|f''\left(\frac{a+b}{2}\right)\right| \end{aligned} \tag{1.6}$$

where β is Euler Beta function.

We will give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further this paper.

Definition 2. *Let $f \in L[a,b]$. The Reimann-Liouville integrals $J_{a^+}^{\alpha} f(x)$ and $J_{b^-}^{\alpha} f(x)$ of order $\alpha > 0$ with $\alpha \geq 0$ are defined by*

$$J_{a^+}^{\alpha}f(x)=\frac{1}{\Gamma(\alpha)}\int_a^x (x-t)^{\alpha-1}f(t)dt, x>\alpha$$

and

$$J_{b^-}^{\alpha}f(x)=\frac{1}{\Gamma(\alpha)}\int_x^b (t-x)^{\alpha-1}f(t)dt, x<b$$

respectively. Where $\Gamma(\alpha)=\int_0^{\infty}e^{-t}u^{\alpha-1}du$ is the Gamma function and $J_{a^+}^0f(x)=J_{b^-}^0f(x)=f(x)$.

In the case of $\alpha=1$ the fractional integral reduces to the classical integral.

For some recent results connected with fractional integral ineqalities, (see [18]-[27]).

In this paper, we establish fractional integral ineqalities of Hermite-Hadamard type for mappings whose derivatives are $s-$convex and concave.

2. Main Results

In order to prove our main theorems we need the following lemma (see [2]).

Lemma 1. *Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a twice differentiable function on I°,the interior of I. Assume that $a,b\in I^{\circ}$ with $a<b$ and $f^{''}\in L[a,b]$, then the following identity for fractional integral with $\alpha>0$ holds:*

$$\begin{aligned}(2.1)\qquad &\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\\ =&\frac{(b-a)^2}{2(\alpha+1)}\int_0^1 t(1-t^{\alpha})\left[f''(ta+(1-t)b)+f''((1-t)a+tb)\right]dt\end{aligned}$$

where $\Gamma(\alpha)=\int_0^{\infty}e^{-t}u^{\alpha-1}du$.

Theorem 6. *Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a twice differentiable function on I° and let $a,b\in I^{\circ}$ with $a<b$ and $f^{''}\in L[a,b]$. If $|f''|$ is $s-$convex in the second sense on I for some fixed $s\in(0,1]$, then the following inequality for fractional integrals with $\alpha>0$ holds:*

$$\begin{aligned}(2.2)\qquad &\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\right|\\ \leq&\frac{(b-a)^2}{2(\alpha+1)}\left[\frac{\alpha}{(s+2)(\alpha+s+2)}+\beta(2,s+1)-\beta(\alpha+2,s+1)\right]\left[|f''(a)|+|f''(b)|\right]\end{aligned}$$

where β is Euler Beta function.

Proof. From Lemma 1 since $|f''|$ is $s-$convex in the second sense on I, we have

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^{+}}^{\alpha}f(b)+J_{b^{-}}^{\alpha}f(a)\right]\right| \\
\leq\ & \frac{(b-a)^{2}}{2(\alpha+1)}\int_{0}^{1}\left|t\left(1-t^{\alpha}\right)\right|\left[\left|f''\left(ta+(1-t)b\right)\right|+\left|f''\left((1-t)a+tb\right)\right|\right]dt \\
\leq\ & \frac{(b-a)^{2}}{2(\alpha+1)}\left[\int_{0}^{1}t(1-t^{\alpha})\left[t^{s}\left|f''(a)\right|+(1-t)^{s}\left|f''(b)\right|\right]dt+\int_{0}^{1}t(1-t^{\alpha})\left[(1-t)^{s}\left|f''(a)\right|+t^{s}\left|f''(b)\right|\right]dt\right] \\
=\ & \frac{(b-a)^{2}}{2(\alpha+1)}\left[\int_{0}^{1}t^{s+1}\left(1-t^{\alpha}\right)dt+\int_{0}^{1}t(1-t^{\alpha})(1-t)^{s}dt\right]\left[\left|f''(a)\right|+\left|f''(b)\right|\right] \\
=\ & \frac{(b-a)^{2}}{2(\alpha+1)}\left[\frac{\alpha}{(s+2)(\alpha+s+2)}+\beta(2,s+1)-\beta(\alpha+2,s+1)\right]\left[\left|f''(a)\right|+\left|f''(b)\right|\right]
\end{aligned}$$

where we used the fact that

$$\int_{0}^{1}t^{s+1}\left(1-t^{\alpha}\right)dt=\frac{\alpha}{(s+2)(\alpha+s+2)}$$

and

$$\int_{0}^{1}t(1-t^{\alpha})(1-t)^{s}dt=\beta(2,s+1)-\beta(\alpha+2,s+1)$$

which completes the proof. □

Remark 1. *In Theorem 6 if we choose $s=1$ then (2.2) reduces the ineqality (1.3) of Theorem 2.*

Theorem 7. *Let $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ be a twice differentiable function on I°. Suppose that $a,b\in I^{\circ}$ with $a<b$ and $f^{''}\in L[a,b]$. If $|f''|^{q}$ is $s-$convex in the second sense on I for some fixed $s\in(0,1]$, $p,q>1$ then the following inequality for fractional integrals holds:*

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^{+}}^{\alpha}f(b)+J_{b^{-}}^{\alpha}f(a)\right]\right| \\
\leq\ & \frac{(b-a)^{2}}{\alpha+1}\beta^{\frac{1}{p}}\left(p+1,\alpha p+1\right)\left[\frac{\left|f''(a)\right|^{q}+\left|f''(b)\right|^{q}}{s+1}\right]^{\frac{1}{q}}
\end{aligned}\tag{2.3}$$

where β is Euler Beta function and $\frac{1}{p}+\frac{1}{q}=1$.

Proof. From Lemma 1,using the well known Hölder inequality and $|f''|^q$ is $s-$convex in the second sense on I, we have

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^{+}}^{\alpha}f(b)+J_{b^{-}}^{\alpha}f(a)\right]\right| \\
\leq\ & \frac{(b-a)^{2}}{2(\alpha+1)}\int_{0}^{1}\left|t\left(1-t^{\alpha}\right)\right|\left[\left|f''\left(ta+(1-t)b\right)\right|+\left|f''\left((1-t)a+tb\right)\right|\right]dt \\
\leq\ & \frac{(b-a)^{2}}{2(\alpha+1)}\left(\int_{0}^{1}t^{p}\left(1-t^{\alpha}\right)^{p}dt\right)^{1-\frac{1}{q}}\left[\left(\int_{0}^{1}|f''(ta+(1-t)b)|^{q}dt\right)^{\frac{1}{q}}+\left(\int_{0}^{1}|f''((1-t)a+tb)|^{q}dt\right)^{\frac{1}{q}}\right] \\
\leq\ & \frac{(b-a)^{2}}{2(\alpha+1)}\left(\int_{0}^{1}t^{p}\left(1-t^{\alpha}\right)^{p}dt\right)^{1-\frac{1}{q}}\left[\begin{array}{c}\left(\int_{0}^{1}\left(t^{s}|f''(a)|^{q}+(1-t)^{s}|f''(b)|^{q}\right)dt\right)^{\frac{1}{q}} \\ +\left(\int_{0}^{1}\left((1-t)^{s}|f''(a)|^{q}+t^{s}|f''(b)|^{q}\right)dt\right)^{\frac{1}{q}}\end{array}\right] \\
=\ & \frac{(b-a)^{2}}{2(\alpha+1)}\left(\int_{0}^{1}t^{p}\left(1-t^{\alpha}\right)^{p}dt\right)^{1-\frac{1}{q}}\left[\begin{array}{c}\left(|f''(a)|^{q}\frac{1}{s+1}+|f''(b)|^{q}\frac{1}{s+1}\right)^{\frac{1}{q}} \\ +\left(|f''(a)|^{q}\frac{1}{s+1}+|f''(b)|^{q}\frac{1}{s+1}\right)^{\frac{1}{q}}\end{array}\right] \\
\leq\ & \frac{(b-a)^{2}}{\alpha+1}\beta^{\frac{1}{p}}\left(p+1,\alpha p+1\right)\left[\frac{|f''(a)|^{q}+|f''(b)|^{q}}{s+1}\right]^{\frac{1}{q}}
\end{aligned}$$

where we used the fact that

$$\int_{0}^{1}t^{s}dt=\int_{0}^{1}(1-t)^{s}dt=\frac{1}{s+1}$$

and

$$\int_{0}^{1}t^{p}\left(1-t^{\alpha}\right)^{p}dt\leq\int_{0}^{1}t^{p}(1-t)^{\alpha p}dt=\beta^{\frac{1}{p}}\left(p+1,\alpha p+1\right)$$

which completes the proof. □

Remark 2. *In Theorem 7 if we choose* $s=1$ *then (2.3) reduces the ineqality (1.4) of Theorem 3.*

Theorem 8. *Let* $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ *be a twice differentiable function on* I°*.Suppose that* $a,b\in I^{\circ}$ *with* $a<b$ *and* $f^{''}\in L[a,b]$*. If* $|f''|^{q}$ *is* $s-$*convex in the second sense on* I *for some fixed* $s\in(0,1]$ *and* $q\geq1$ *then the following inequality for fractional integrals holds:*

$$\begin{aligned}
(2.4)\quad &\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^{+}}^{\alpha}f(b)+J_{b^{-}}^{\alpha}f(a)\right]\right| \\
\leq\ & \frac{\alpha(b-a)^{2}}{4(\alpha+1)(\alpha+2)} \\
&\times\left[\begin{array}{c}\left(|f''(a)|^{q}\frac{2\alpha+4}{(s+2)(\alpha+s+2)}+|f''(b)|^{q}\left[\beta(2s+1)-\beta(\alpha+2,s+1)\right]\frac{2\alpha+4}{\alpha}\right)^{\frac{1}{q}} \\ \left(|f''(a)|^{q}\left[\beta(2s+1)-\beta(\alpha+2,s+1)\right]\frac{2\alpha+4}{\alpha}+|f''(b)|^{q}\frac{2\alpha+4}{(s+2)(\alpha+s+2)}\right)^{\frac{1}{q}}\end{array}\right]
\end{aligned}$$

Proof. From Lemma 1, using power mean ineqality and $|f''|^q$ is $s-$convex in the second sense on I we have

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\right| \\
\leq\ & \frac{(b-a)^2}{2(\alpha+1)}\int_0^1 |t(1-t^{\alpha})|\left[|f''(ta+(1-t)b)|+|f''((1-t)a+tb)|\right]dt \\
\leq\ & \frac{(b-a)^2}{2(\alpha+1)}\left(\int_0^1 t(1-t^{\alpha})\,dt\right)^{1-\frac{1}{q}}\left[\begin{array}{c}\left(\int_0^1 t(1-t^{\alpha})\,|f''(ta+(1-t)b)|^q\,dt\right)^{\frac{1}{q}} \\ +\left(\int_0^1 t(1-t^{\alpha})\,|f''((1-t)a+tb)|^q\,dt\right)^{\frac{1}{q}}\end{array}\right] \\
\leq\ & \frac{(b-a)^2}{2(\alpha+1)}\left(\int_0^1 t(1-t^{\alpha})\,dt\right)^{1-\frac{1}{q}}\left[\begin{array}{c}\left(\int_0^1 \left[t^{s+1}(1-t^{\alpha})\,|f''(a)|^q+t(1-t^{\alpha})(1-t)^s\,|f''(b)|^q\right]dt\right)^{\frac{1}{q}} \\ +\left(\int_0^1 t(1-t^{\alpha})(1-t)^s\,|f''(a)|^q+t^{s+1}(1-t^{\alpha})\,|f''(b)|^q\,dt\right)^{\frac{1}{q}}\end{array}\right] \\
=\ & \frac{(b-a)^2}{2(\alpha+1)}\left(\frac{\alpha}{2(\alpha+2)}\right)^{1-\frac{1}{q}} \\
& \times\left[\begin{array}{c}\left(|f''(a)|^q\,\frac{\alpha}{(s+2)(\alpha+s+2)}+|f''(b)|^q\left[\beta(2s+1)-\beta(\alpha+2,s+1)\right]\right)^{\frac{1}{q}} \\ +\left(|f''(a)|^q\left[\beta(2s+1)-\beta(\alpha+2,s+1)\right]+|f''(b)|^q+|f''(b)|^q\,\frac{\alpha}{(s+2)(\alpha+s+2)}\right)^{\frac{1}{q}}\end{array}\right] \\
\leq\ & \frac{\alpha(b-a)^2}{4(\alpha+1)(\alpha+2)} \\
& \times\left[\begin{array}{c}\left(|f''(a)|^q\,\frac{2\alpha+4}{(s+2)(\alpha+s+2)}+|f''(b)|^q\left[\beta(2s+1)-\beta(\alpha+2,s+1)\right]\frac{2\alpha+4}{\alpha}\right)^{\frac{1}{q}} \\ \left(|f''(a)|^q\left[\beta(2s+1)-\beta(\alpha+2,s+1)\right]\frac{2\alpha+4}{\alpha}+|f''(b)|^q\,\frac{2\alpha+4}{(s+2)(\alpha+s+2)}\right)^{\frac{1}{q}}\end{array}\right]
\end{aligned}$$

where we used the fact that

$$\int_0^1 t^{s+1}(1-t^{\alpha})\,dt=\frac{\alpha}{(s+2)(\alpha+s+2)}$$

and

$$\int_0^1 t(1-t^{\alpha})(1-t)^s\,dt=\beta(2s+1)-\beta(\alpha+2,s+1)$$

which completes the proof. □

Remark 3. *In Theorem (8) if we choose $s=1$ then(2.4) reduces the ineqality (1.5) of Theorem (4).*

The following result holds for $s-$concavity.

Theorem 9. *Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a twice differentiable function on I°.Suppose that $a,b\in I^{\circ}$ with $a<b$ and $f^{''}\in L[a,b]$. If $|f''|^q$ is $s-$concave in the second sense on I for some fixed $s\in(0,1]$ and $p,q>1$ then the following inequality for fractional integrals holds:*

$$\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\right| \tag{2.5}$$

$$\leq \frac{(b-a)^2}{\alpha+1}\beta^{\frac{1}{p}}(p+1,\alpha p+1)\,2^{\frac{s-1}{q}}\left|f''\left(\frac{a+b}{2}\right)\right|$$

where $\frac{1}{p}+\frac{1}{q}=1$ *and* β *is* β *is Euler Beta function*

Proof. From Lemma 1 and using the Hölder ineqality we have

$$\begin{aligned} & \left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\right| \\ \leq & \frac{(b-a)^2}{2(\alpha+1)}\int_0^1 \left|t\left(1-t^{\alpha}\right)\right|\left[\left|f''\left(ta+(1-t)b\right)\right|+\left|f''\left((1-t)a+tb\right)\right|\right]dt \\ \leq & \frac{(b-a)^2}{2(\alpha+1)}\left(\int_0^1 t^p\left(1-t^{\alpha}\right)^p dt\right)^{\frac{1}{p}}\left[\left(\int_0^1 |f''(ta+(1-t)b)|^q dt\right)^{\frac{1}{q}}+\left(\int_0^1 |f''((1-t)a+tb)|^q dt\right)^{\frac{1}{q}}\right] \\ \leq & \frac{(b-a)^2}{2(\alpha+1)}\left(\int_0^1 t^p\left(1-t^{\alpha}\right)^p dt\right)^{\frac{1}{p}} \\ & \times\left[\begin{array}{c} \left(\int_0^1\left(t^s|f''(a)|^q+(1-t)^s|f''(b)|^q\right)dt\right)^{\frac{1}{q}} \\ +\left(\int_0^1\left((1-t)^s|f''(a)|^q+t^s|f''(b)|^q\right)dt\right)^{\frac{1}{q}} \end{array}\right] \end{aligned} \tag{2.6}$$

Since $|f''|^q$ is $s-$concave using ineqality (1.2) we get (see [1])

$$\int_0^1 |f''(ta+(1-t)b)|^q\,dt \leq 2^{s-1}\left|f''\left(\frac{a+b}{2}\right)\right|^q \tag{2.7}$$

and

$$\int_0^1 |f''((1-t)a+tb)|^q\,dt \leq 2^{s-1}\left|f''\left(\frac{b+a}{2}\right)\right|^q \tag{2.8}$$

Using (2.7) and (2.8) in (2.6), we have

$$\begin{aligned} & \left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right]\right| \\ \leq & \frac{(b-a)^2}{\alpha+1}\beta^{\frac{1}{p}}\left(p+1,\alpha p+1\right)2^{\frac{s-1}{q}}\left|f''\left(\frac{a+b}{2}\right)\right| \end{aligned}$$

which completes the proof. □

Remark 4. *In theorem (9) if we choose* $s=1$ *then (2.5) reduces ineqality (1.6) of theorem 5.*

References

[1] M. Alomari, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives are s-convex in the second sense, Appl. Math. Lett. 23 (2010) 1071-1076.

[2] S.S.Dragomir,M.I.Bhatti and M.Iqbal Some new fractional integral Hermite-Hadamard type inequalities, RGMIA Res. Rep. Coll., 16 (2013), Article 2.

[3] S.S.Dragomir,S.Fitzpatrik,The Hadamard's inequality for s-convex functions in the second sense, Demonstratio Math 32 (4) (1999) 687-696.

[4] H. Hudzik, L. Maligranda, Some remarks on s-convex functions, Aequationes Math. 48 (1994) 100–111.

[5] M. Alomari, M. Darus, On the Hadamard's inequality for log-convex functions on the coordinates, J. Inequal. Appl. 2009 (2009) 13. Article ID 283147.

[6] A.G. Azpeitia, Convex functions and the Hadamard inequality, Revista Colombiana Mat. 28 (1994) 7–12.

[7] M.K. Bakula, M.E. Özdemir, J. Pečarić, Hadamard tpye inequalities for m-convex and (α,m)-convex functions, J. Ineq. Pure Appl. Math. 9 (4) (2008) Art. 96.
[8] M.K. Bakula, J. Pečarić, Note on some Hadamard-type inequalities, J. Ineq. Pure Appl. Math. 5 (3) (2004) Article 74.
[9] S.S. Dragomir, C.E.M. Pearce, Selected topics on Hermite–Hadamard inequalities and applications, RGMIA Monographs, Victoria University, 2000.
[10] S.S. Dragomir, R.P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula,Appl. Math. lett. 11 (5) (1998) 91–95.
[11] S.S. Dragomir, On some new inequalities of Hermite–Hadamard type for m-convex functions, Tamkang J. Math. 3 (1) (2002).
[12] P.M. Gill, C.E.M. Pearce, J. Pečarić, Hadamard's inequality for r-convex functions, J. Math. Anal. Appl. 215 (2) (1997) 461–470.
[13] U.S. Kırmacı, M.K. Bakula, M.E. Özdemir, J. Pečarić, Hadamard-tpye inequalities for s-convex functions, Appl. Math. Comput. 193 (2007) 26–35.
[14] M.E. Özdemir, M. Avci, E. Set, On some inequalities of Hermite–Hadamard type via m-convexity, Appl. Math. Lett. 23 (9) (2010) 1065–1070.
[15] E. Set, M.E. Özdemir, S.S. Dragomir, On the Hermite–Hadamard inequality and other integral inequalities involving two functions, J. Inequal. Appl.(2010) 9. Article ID 148102.
[16] E. Set, M.E. Özdemir, S.S. Dragomir, On Hadamard-type inequalities involving several kinds of convexity, J. Inequal. Appl. (2010) 12. Article ID 286845.
[17] J.E. Pečarić, F. Proschan, Y.L. Tong, Convex Functions, Partial Orderings and Statistical Applications, Academic Press, Boston, 1992.
[18] G. Anastassiou, M.R. Hooshmandasl, A. Ghasemi, F. Moftakharzadeh, Montogomery identities for fractional integrals and related fractional inequalities J. Ineq. Pure Appl. Math. 10 (4) (2009) Art 97.
[19] S. Belarbi, Z. Dahmani, On some new fractional integral inequalities, J. Ineq. Pure Appl. Math. 10 (3) (2009) Art. 86.
[20] Z. Dahmani, New inequalities in fractional integrals, Int. J. Nonlinear Sci. 9 (4) (2010) 493–497.
[21] Z. Dahmani, On Minkowski and Hermite–Hadamard integral inequalities via fractional integration, Ann. Funct. Anal. 1 (1) (2010) 51–58.
[22] Z. Dahmani, L. Tabharit, S. Taf, Some fractional integral inequalities, Nonlinear. Sci. Lett. A 1 (2) (2010) 155–160.
[23] Z. Dahmani, L. Tabharit, S. Taf, New generalizations of Gruss inequality usin Riemann–Liouville fractional integrals, Bull. Math. Anal. Appl. 2 (3) (2010) 93–99.
[24] M.Z. Sarikaya, H. Ogunmez, On new inequalities via Riemann–Liouville fractional integration, Abstract and Applied Analysis, Volume 2012, Article ID 428983, 10 pages, doi:10.1155/2012/428983.
[25] M.Z. Sarikaya and H. Yaldiz, On weighted Montogomery identities for Riemann-Liouville fractional integrals, Konuralp Journal of Mathematics, 1 (1) (2013) 48-53.
[26] M.Z. Sarikaya, E. Set, H. Yaldiz and N. Basak, Hermite -Hadamard's inequalities for fractional integrals and related fractional inequalities, Mathematical and Computer Modelling, doi:10.1016/j.mcm.2011.12.048, in press.
[27] E. Set, New inequalities of Ostrowski type for mappings whose derivatives are s-convex in the second sense via fractional integrals, Comp. Math. Appl., 63(7) (2012), 1147-1154.

[▲]Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-TURKEY
E-mail address: `erhanset@yahoo.com`

[★]Atatürk University, K.K. Education Faculty, Department of Mathematics, 25240, Campus, Erzurum, Turkey
E-mail address: `emos@atauni.edu.tr`

[■]Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-TURKEY
E-mail address: `sarikayamz@gmail.com`

[♣]Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-TURKEY
E-mail address: `filinz_41@hotmail.com`